




031606

17712 U.S.PTO

ELECTRON MICROGRAPHS OF ULTRATHIN SECTIONS OF  
LYMPHOCYTES OF A HEALTHY ADULT DONOR INFECTED WITH THE  
RETROVIRUS ISOLATED FROM A HAEMOPHILIAC PATIENT WITH AIDS (IDAV2 )



NOTE THE NUMEROUS BUDDING WITH PROJECTIONS AT THEIR SURFACE  
AND A MATURE PARTICLE WITH A SMALL CORE.


**FIG. 1**

BEST AVAILABLE COPY

## Replacement Sheet

17712 U.S. PTO  
031606


ULTRASTRUCTURAL COMPARISON BETWEEN HTLV AND LAV.



HTLV PARTICLES PRODUCED BY C10 MJ<sub>2</sub> CELL LINE.  
NOTE THE LARGE CORE OF MATURE PARTICLES AND A TYPICAL BUDDING.

### **FIG. 2A**

ULTRASTRUCTURAL COMPARISON BETWEEN HTLV AND LAV.



LAV MATURE PARTICLES WITH DENSE CORE AND  
ONE BUDDING PRODUCED BY INFECTED LYMPHOCYTES.

### **FIG. 2B**

BEST AVAILABLE COPY

Replacement Sheet

BEST AVAILABLE COPY

# FIG. 3

INTRACELLULAR VESICLE IN LAV - INFECTED LYMPHOCYTES  
ARROW INDICATES A BUDDING PARTICLE



17712 U.S. PTO




081606

Replacement Sheet

17712 U.S. PTO  
031606

ELECTROPHORESIS AND AUTORADIOGRAM OF  
<sup>35</sup>METHIONINE LABELLED LAV.




AUTORADIOGRAM OF THIS GEL, WITH, ON THE LEFT,  
MOLECULAR WEIGHT MARKERS IN KILODALTONS. NOTE THAT  
THE p25 PROTEIN COINCIDES WITH THE PEAK OF LABELLED  
VIRUS AND THAT OF REVERSE TRANSCRIPTASE ACTIVITY  
(NOT REPRESENTED)

**FIG. 4A**

BEST AVAILABLE COPY

# Replacement Sheet



2ND PANEL: BANDING OF LAV IN A NYCODENZ GRADIENT. INFECTED LYMPHOCYTES FROM A HEALTHY DONOR WERE LABELLED FOR 18 h IN THE PRESENCE OF <sup>35</sup>S-METHIONINE, AS DESCRIBED IN [1]. VIRUS WAS PRECIPITATED FROM THE CLARIFIED SUPERNATANT WITH 10% PEG 6000 OVERNIGHT AT 4°C AND THE PELLET WAS RESUSPENDED IN 0.5ml OF NTE BUFFER (0.1M NaCl, 0.01M TRIS, 0.001M EDTA, pH 7.4). IT WAS THEN Banded TO EQUILIBRIUM IN A LINEAR NYCODENZ (NYEGAAND, OSLO, 5.35%) GRADIENT IN A SW56 ROTOR FOR 3 h AT 45,000 RPM, 2°C. ALIQUOTS OF THE COLLECTED FRACTION WERE ASSAYED FOR RT ACTIVITY (10 ul), RADIOACTIVITY (20 ul; THICK LINE), AND 40 ul WERE ELECTROPHORESSED ON A POLYACRYLAMIDE GEL (12.5%) UNDER DENATURING CONDITIONS. DENSITY OF RETROVIRUSES IN NYCODENZ GRADIENTS (LAV OR MuLV) WAS VERY LOW (AROUND 1.10).

**FIG. 4B**

BEST AVAILABLE COPY